



Belgrave Medical Centre

THE SPECIALIST HAIR-LOSS CENTRE

LIQUID TREATMENT APPLICATION

Your application should be as follows:

- 1) Apply lotion to dry scalp either after shampooing and blow drying or without shampooing.
- 2) Make a parting across the scalp from frontal hairline to crown starting at the mid eyebrow point.
- 3) Put the syringe dropper into the bottle and draw the lotion up into the syringe to the 1.0ml marker. This fills the whole syringe.
- 4) Holding the hair flat with one hand, place a few drops of lotion along the whole of the parting. Quickly dispense the product across the skin with the fingers to help absorption.
- 5) Make another parting approximately 1cm up from the first parting (towards top of head). Repeat application.
- 6) Continue to make partings approximately 1 cm apart across the top of the head to mid eyebrow point on the other side of head.
- 7) Any surplus product can then be used throughout the area.
- 8) This effectively covers the entire area of the upper scalp.
- 9) Wash hands thoroughly.
- 10) Repeat process at bedtime.

NOTE:

- 1) You only need 1.0ml to cover the entire upper scalp.
- 2) You do not need to refill the syringe for each parting.
- 3) Do not apply near a naked flame.

MINOXIDIL IN CONJUNCTION WITH OESTRADIOL OR ESTRADEX

- 1) If using minoxidil only apply 1ml in the morning and then repeat application before bedtime.
- 2) If using minoxidil in conjunction with estradiol or estradex apply 1ml of oestradiol or estradex in the morning and 1ml of minoxidil before bedtime using the same application method.
- 3) If using twice daily oestradiol or estradex without minoxidil apply 1ml of either solution in the morning and repeat application at bedtime using the same application method.

Diagram:

